Your procedure will be performed by Doctor: _____

Address: _____

Date of procedure:_____Arrive at:_____AM/PM

Comments: _____

SUPREP® BOWEL PREP KIT

(sodium sulfate, potassium sulfate and magnesium sulfate) Oral Solution

For Adults

(17.5g/3.13g/1.6g) per 6 ounces



On the day *before* your procedure…

What You **CAN** do:

- You may have a light breakfast or have clear liquids ONLY; please have nothing for dinner

What You **CANNOT** do:

- **DO NOT** drink milk
- **DO NOT** eat or drink anything colored red or purple
- **DO NOT** drink alcoholic beverages

Any of the following liquids are OK to drink:

- Water
- Strained fruit juices (without pulp) including apple, orange, white grape, or white cranberry
- Limeade or lemonade
- Coffee or tea (**DO NOT** use any dairy or non-dairy creamer)
- Chicken broth
- Gelatin desserts without added fruit or topping **(NO RED OR PURPLE)**

Note:

- SUPREP Bowel Prep Kit is an osmotic laxative indicated for cleansing of the colon as a preparation for colonoscopy in adult and pediatric patients 12 years of age and older
- Be sure to tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. SUPREP Bowel Prep Kit may affect how other medicines work
- Medication taken by mouth may not be absorbed properly when taken within 1 hour before the start of each dose of SUPREP Bowel Prep Kit
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these medications at least 2 hours before and not less than 6 hours after administration of SUPREP Bowel Prep Kit
- The most common adverse reactions in adults (>2%) after administration of SUPREP Bowel Prep Kit were overall discomfort, abdominal distention, abdominal pain, nausea, and vomiting
- If you have any questions about taking SUPREP Bowel Prep Kit, call your doctor


To learn more about this product, please call **1-800-874-6756** or visit our Web site at **www.suprepkit.com**

SUPREP Bowel Prep Kit is a split-dose (2-day) regimen. Both 6-ounce bottles are required for a complete prep.

FIRST DOSE: Begin Step 1 at _____PM the evening before your procedure and proceed as shown below:


You must complete Steps 1 through 4 using one (1) 6-ounce bottle before going to bed:

STEP 1



Pour **ONE** (1) 6-ounce bottle of SUPREP liquid into the mixing container.


STEP 2



Add cool drinking water to the 16-ounce line on the container and mix.


***NOTE:** Dilute the solution concentrate as directed prior to use.*

STEP 3



Drink **ALL** the liquid in the container.

STEP 4



You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

SECOND DOSE: Begin Step 1 at _____AM the morning of your procedure and proceed as shown above:

For this dose, repeat Steps 1 through 4 shown above using the other 6-ounce bottle.

***NOTE:** You **must** finish drinking the final glass of water at least 2 hours, or as directed, before your colonoscopy.*

Please read the full Prescribing Information and Medication Guide in the kit.

Provided as an educational service from Braintree A PART OF SEBELA PHARMACEUTICALS®

©2021 Braintree Laboratories, Inc.

012-271-v1 March 2021